

 <b>Lakeridge Health</b>	<b>Neonatal and Paediatric Intubation and LMA Insertion – Medical Directive</b>	
	Manual: Medical Directives & Delegated Controlled Acts	Original Date: 17FEB2009
	Section: Paediatrics and Neonatology	Version Date: 24JUN2025
	Document Owner: Critical Care	Next Review Date: 24JUN2027
	Approved by: Women’s and Children’s Council, Interprofessional Collaboration Committee, Operations Committee, Medical Advisory Committee	
Cross Reference to: Endotracheal Intubation Policy and Procedures		
<p>A printed copy of this document may not reflect the current, electronic version on Lakeridge Health’s Intranet, ‘OASIS’. Any copies of this document appearing in paper form should ALWAYS be checked against the electronic version prior to use. Contact policies@lh.ca for version history.</p>		

## Authorizing Prescribers

- Lakeridge Health Oshawa (LHO) Emergency Physicians, Paediatricians, Neonatologists, Anesthesiologists
- Lakeridge Health Ajax-Pickering (LHAP) Emergency Physicians, Paediatricians, Neonatologists, Anesthesiologists
- Lakeridge Health Bowmanville (LHB), Lakeridge Health Port Perry (LHPP) Emergency Physicians, Anesthesiologists

## Authorized to Whom

Registered Respiratory Therapists (RRT) and Anesthesia Assistants (AA) employed within Lakeridge Health who have reviewed the affiliated policy and demonstrated competency supported by the Clinical Practice Leader (CPL) or delegate. They must also be in good standing with the College of Respiratory Therapists of Ontario and perform as authorized to them under Controlled Act #2 - “intubation beyond the point in the nasal passages where they normally narrow or beyond the larynx.”

**Co-implementer:** Medical Radiation Technologists - Radiography (MRT(R))

## Patient Description/Population

Any person that is or appears to be 12 years old and younger within Lakeridge Health (LH) who meets procedure-specific indications as per the order table.

## Order and/or Procedure

These procedures are not presented in sequential order and any one of or combination of the procedures below may be performed by an authorized health care professional as time permits until either an authorized prescriber or appropriate emergency response personnel are present.

### A. Insertion of an Endotracheal Tube (ETT)

---

This material has been prepared solely for the use at Lakeridge Health. Lakeridge Health accepts no responsibility for use of this material by any person or organization not associated with Lakeridge Health. No part of this document may be reproduced in any form for publication without the permission of Lakeridge Health.

## B. Insertion of a Laryngeal Mask Airway (LMA)

### Indications to the Implementation of the Directive

- Any patient with procedure-specific indications as per [Order Table](#)

### Contraindications to the Implementation of the Directive

This medical directive must not be implemented in the following circumstances:

- In a known or suspected cervical spine injury or instability
- The patient, their guardian or Substitute Decision Maker (SDM) does not consent to resuscitation. It is either clearly expressed and/or documented in the patient's medical record
- The RRT or AA has not maintained competency in neonatal, paediatric or adult intubation
- In an urgent or semi-urgent upper airway obstruction (i.e., croup, acute epiglottitis, congenital anomalies, oropharyngeal tumors)
- Active hemorrhaging from the oral or nasopharynx

### Consent

The RRT or AA implementing the directive must obtain consent if the patient or Substitute Decision Maker (SDM) is capable of providing it. In an emergency situation, if the patient is not capable of providing consent, the RRT or AA may administer treatment without consent if, in his or her opinion, all of the following are true:

- The patient is incapable with respect to the treatment
- The patient is experiencing severe suffering or is at risk if the treatment is not administered promptly, or suffering serious bodily harm
- It is not reasonably possible to obtain a consent or refusal on the patient's behalf, or the delay required to do so will prolong the suffering than the patient is experiencing or will put the patient at risk of suffering serious bodily harm.

### Documentation Requirements

In addition to standard documentation practices, the RHCP implementing this medical directive must document in the order section of the chart (if documenting electronically, document in assessment form or patient note) the following:

- The procedure performed on the patient
- The name of this medical directive
- The name of the implementer
- Legible signature of implementer including credentials (unless documenting electronically)
- Date and time (unless documenting electronically)

### Review/Evaluation Process

This directive will be reviewed by Critical Care Council every 2 years.

## **Reference**

Jannu, A., Shekar, A., Balakrishna, R., Sudarshan, H., Veena, G. C., & Bhuvaneshwari, S. (2017). Advantages, Disadvantages, Indications, Contraindications and Surgical Technique of Laryngeal Airway Mask. *Archives of craniofacial surgery*, 18(4), 223–229. <https://doi.org/10.7181/acfs.2017.18.4.223>

\*\*\*This table must **not** be used independently apart from the Medical Directive\*\*\*

**Order Table Form**

Order(s)	Indications	Contraindications	Notes
Insertion of Endotracheal Tube  Portable chest x-ray post-ETT insertion to confirm placement	<ul style="list-style-type: none"> <li>• Any patient in a cardiac or respiratory arrest (impending or actual)</li> <li>• In patients who have decreased LOC and are either unable to protect their airway or maintain ventilation</li> <li>• Accidental displacement of an endotracheal tube in which the patient is unable to maintain/protect airway, or maintain adequate oxygenation and/or ventilation.</li> </ul>	<ul style="list-style-type: none"> <li>• In a known or suspected cervical spine injury or instability</li> <li>• The patient, their guardian or Substitute Decision Maker (SDM) does not consent to resuscitation. It is either clearly expressed and/or documented in the patient's medical record</li> <li>• The RRT or AA has not maintained competency in neonatal, paediatric or adult intubation</li> <li>• In an urgent or semi-urgent upper airway obstruction (i.e. croup, acute epiglottitis, congenital anomalies, oropharyngeal tumors)</li> <li>• Active hemorrhaging from the oral or nasopharynx</li> </ul>	<ul style="list-style-type: none"> <li>• Pre-oxygenate the patient as needed</li> <li>• Suction prior to inserting ETT or LMA</li> <li>• LMA can be inserted during CPR</li> <li>• Ensure MRP is aware of the treatment initiated</li> </ul>
Insertion of LMA			