



**Lakeridge
Health**

Nuclear Medicine Requisition

Lakeridge Health Oshawa, 1 Hospital Court, Oshawa ON L1G 2B9
Telephone: 905-576-8711 Extension 3345 Fax: 905-721-4762

To book a scan, call Central Booking Office – Telephone (8am to 5pm): 905-721-4717

Requisition must be faxed by Dr's Office or brought by patient to Nuclear Medicine Dept.

Patient's Name: _____ Date of Birth _____ Phone: _____
(DDMMYYYYY)

Appointment Date: _____ Time: _____
(DDMMYYYYY)

**Bring list of medications
and Health Card**

Referring Physician: _____
(PRINT Name) (Signature)

Send copy of report to: _____

Bone Scan (flow, pool, SPECT if needed)

- ☐ Total body
- ☐ Single Site
- ☐ Facial/head and neck SPECT Scan
- ☐ Biliary scan with flow
- ☐ Biliary scan with flow and CCK
- ☐ Bone Mineral Densitometry
- ☐ Brain Perfusion SPECT with flow
- ☐ Gallium Scan
- ☐ Gastric Emptying Study
- ☐ GI Bleed (RBC Label with flow)
- ☐ Meckel's Scan
- ☐ I-131 Metastatic Survey
- ☐ Liver/Spleen Scan with flow
- ☐ Liver/RBC Scan with flow and BPs (for hemangioma)

Lung Scan

- ☐ Ventilation/Perfusion
- ☐ Quantitative – for pre-op assessment

Lymphangiogram (Sentinal lymph node)

- ☐ Breast
- ☐ Melanoma
- ☐ Other _____

Myocardial Perfusion Scan with EF

- ☐ Exercise Treadmill Stress
Weight _____ Height _____
- ☐ Pharmacological Stress (i.e. Persantine)
Weight _____ Height _____
- ☐ Myocardial Viability Scan (Thallium)
- ☐ MUGA Rest
- ☐ MUGA Rest and Exercise
- ☐ Parathyroid
- ☐ Renal Scan (with GFR)
 - ☐ With ACE inhibitor
 - ☐ With Lasix
- ☐ Renal DMSA Scan for renal scar assessment
- ☐ Salivary Scan with flow
- ☐ Testicular Scan with flow

Thyroid

- ☐ Uptake and Scan
- ☐ Scan only (99m Tc)
- ☐ WBC Scan (with Colloid Scan if applicable)

Clinical Information: (Please be specific) _____

For Technologist Use Only:

Are you pregnant? ☐ Yes ☐ No
Are you breastfeeding? ☐ Yes ☐ No

Patient's Initials: _____

Start Time: _____ End Time: _____

- ☐ Order Amended
- ☐ Paperwork Scanned
- ☐ Sent to PACS
- ☐ Previous Study Sent
- ☐ Taken
- ☐ Sent to Med Image
- ☐ Verified

<input type="checkbox"/> I.V.	<input type="checkbox"/> I.M.	<input type="checkbox"/> P.O.
Pharmaceutical:		
Site:	Time:	
Comments:		
MRT (NM):		



Patient Preparations	Estimated Time in Nuclear Medicine
Note: For studies not listed here, there is no prep unless indicated by your doctor. Most of our studies take at least 1 hour to complete.	
Biliary Scan (with or without CCK): No food or drink 4 hrs prior to test. No morphine 4 hours prior to test.	1.5 to 2 hours
Bone Mineral Density: No barium or Gastrografin for 7 days prior to test.	30 minutes
Bone Scan: No prep required.	2 visits required: first visit (20–30 minutes) for injection ; return 2–4 hours later for imaging (30–60 minutes). You may leave the department between visits.
Gallium Scan: No prep required	2 visits required: first day for injection (20 minutes); second visit patient returns 48–72 hours later for imaging (45–90 minutes). Sometimes additional imaging is required in the days following the initial imaging.
Gastric Emptying: No food or drink 4 hours prior to test.	4–5 hours
GI Bleed: No barium 3 days prior to test.	1 to 6 hours.
Meckel's Study: No food or drink 4 hours prior to test. No ASA (aspirin) 3 days prior to test. No barium 7 days prior to test. Cimetidine should be prescribed by your Doctor to be taken day before test.	1 hour.
Myocardial Perfusion: No food and no caffeine for 8 hours prior to test. Plus, for pharmacological study, no aminophylline or similar type medications for 2 days prior to test. DIABETIC PATIENTS may have small meal with their insulin or pills (e.g. juice and toast).	99mTC agent (Cardiolite) testing: 5–6 hours if BMI less than 35 If BMI greater than 35, 2 visits are required: * 1st day – 2 hours * 2nd day – 3 hours
Parathyroid Scan: No prep required.	2 visits required: first visit (45 minutes) for injection and image. Return approximately 3 hours later for second image (45 minutes)
RBC Liver Scan: No prep required.	2 visits required: first – 1.5 hours; second – 30 minutes. About 2 hour gap between visits.
Renal Scan: Drink 3–4 glasses of fluid prior to arrival.	1–1.5 hours. DMSA – initial injection. Return 2–4 hours later for imaging.
Salivary Gland: No food or drink 4 hours prior to test.	1 hour.
Thyroid Uptake & Scan: Discontinue thyroid hormone medications for 2 weeks (if referring doctor says it is ok). Discontinue anti-thyroid medications for 5 days. No contrast dye or Gallium 4 weeks prior to test. Stay off kelp or vitamins with iodine for 2 weeks. No food or drink for 4 hours prior to test.	2 visits required 1 day apart: first day (30 minutes) you receive a drink; second day (1 hour) is for imaging.
Thyroid Scan only: Prep same as Thyroid Uptake & Scan above.	1 hour.
WBC Scan: No prep required.	Multiple visits required. First visit (45 minutes) to take blood. Second visit (20 minutes) to have labeled white blood cells re-injected, usually in the afternoon. Third visit (45 minutes) can be same day as other visits or the next day depending on the imaging agent. Additional visits at discretion of Nuclear Medicine Physician.

